

# Herbicide Resistance Management Strategies

*Developed by the CropLife Australia Herbicide Resistance Management Review Group  
and industry researchers*

**Valid as at 15 September 2009**

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## Contents

<b>1. HERBICIDE RESISTANCE .....</b>	<b>3</b>
1.1. Evolution of herbicide resistance.....	3
1.2. Background to herbicide resistance in Australia .....	3
1.3. Current impact on weed management .....	3
1.4. Action by industry and researchers.....	4
<b>2. MODE OF ACTION .....</b>	<b>4</b>
2.1. Mode of action matters!.....	4
2.2. Mode of action labelling in Australia .....	4
<b>3. CHANGES TO MODE OF ACTION GROUPINGS FOR SOME HERBICIDES.</b>	<b>5</b>
<b>4. HERBICIDES ARE GROUPED BY MODE OF ACTION AND RANKED BY RESISTANCE RISK.....</b>	<b>6</b>
<b>5. INTEGRATED WEED MANAGEMENT STRATEGIES.....</b>	<b>7</b>
<b>6. SPECIFIC GUIDELINES FOR GROUP A HERBICIDES.....</b>	<b>9</b>
<b>7. SPECIFIC GUIDELINES FOR GROUP B HERBICIDES.....</b>	<b>10</b>
<b>8. SPECIFIC GUIDELINES FOR GROUP C HERBICIDES.....</b>	<b>12</b>
<b>9. SPECIFIC GUIDELINES FOR GROUP D HERBICIDES.....</b>	<b>14</b>
<b>10. SPECIFIC GUIDELINES FOR GROUP F HERBICIDES .....</b>	<b>15</b>
<b>11. SPECIFIC GUIDELINES FOR GROUP I HERBICIDES .....</b>	<b>16</b>
<b>12. SPECIFIC GUIDELINES FOR GROUP J HERBICIDES .....</b>	<b>17</b>
<b>13. SPECIFIC GUIDELINES FOR GROUP L HERBICIDES .....</b>	<b>18</b>
<b>14. SPECIFIC GUIDELINES FOR GROUP M HERBICIDES .....</b>	<b>20</b>
<b>15. SPECIFIC GUIDELINES FOR GROUP Q HERBICIDES.....</b>	<b>21</b>
<b>16. SPECIFIC GUIDELINES FOR GROUP Z HERBICIDES .....</b>	<b>22</b>

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 1. HERBICIDE RESISTANCE

### 1.1. Evolution of herbicide resistance

Herbicide resistance evolves following the intensive use of herbicides for weed control. In any weed population there are likely to be a small number of individuals that are naturally resistant to herbicides, even before the herbicides are used. When a herbicide is used, these individuals survive and set seed whereas the majority of susceptible plants are killed. Continued use of a herbicide or herbicide group will eventually result in a significant fraction of the weed population with resistance.

There are four main factors that influence the evolution of resistance. These are:

- **The intensity of selection pressure.**  
This refers to how many weeds are killed by the herbicide. For most weeds and herbicides, the number of years of herbicide use is a good measure of selection intensity.
- **The frequency of use of a herbicide or mode of action group.**
- **The frequency of resistance present in untreated populations.**  
If the frequency of resistance risk is relatively high, such as with Group B herbicides, resistance will occur quickly. If the frequency is low, such as with Group M herbicides, resistance will occur more slowly.
- **The biology and density of the weed.**  
Weed species that produce large numbers of seed and have a short seed bank life in the soil will evolve resistance faster than weed species with long seed bank lives. Weed species with greater genetic diversity are more likely to evolve resistance. Resistance is also more likely to be detected in larger weed populations.

### 1.2. Background to herbicide resistance in Australia

Herbicide resistance has developed a strong foothold in Australian agriculture since it was first reported in annual ryegrass in 1982. It has spread and diversified to become a key constraint to crop production in all states generally with a history of intensive herbicide use.

### 1.3. Current impact on weed management

Today, resistance has been confirmed in 34 grass and broadleaf weed species. More worrying still, resistance has now developed to 11 distinctly different herbicide chemical groups. This significantly reduces herbicide options for the grower. Cases of multiple resistance have also been commonly reported where, for example, annual ryegrass proves resistant to two or more chemical groups.

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## **1.4. Action by industry and researchers**

CropLife Australia, with support from the CRC for Australian Weed Management and the Grains Research and Development Corporation (GRDC), introduced a classification system for herbicides enabling farmers and advisers to understand the mode of action grouping. It is mandatory for all herbicide product labels in Australia to carry the designated mode of action group letter in a prominent position. A survey of growers and agronomists (Kondinin, 1998) revealed that 85% of growers are aware of herbicide mode of action groups and consider this important when making buying decisions. This was a good start but resistance management strategies require continual implementation.

## **2. MODE OF ACTION**

### **2.1. Mode of action matters!**

The main reason resistance has developed is because of the repeated and often uninterrupted use of herbicides with the same mode of action. Selection of resistant strains can occur in as little as 3-4 years if no attention is paid to resistance management. Remember that the resistance risk is the same for products having the same mode of action. If you continue to use herbicides with the same mode of action and do not follow a resistance management strategy you are creating future problems for yourself. Mode of action matters.

### **2.2. Mode of action labelling in Australia**

In order to facilitate management of herbicide resistant weeds, all herbicides sold in Australia are grouped by mode of action. The mode of action is indicated by a letter code on the product label. The mode of action labelling is based on the resistance risk of each group of herbicides. Australia was the first country to introduce compulsory mode of action labelling on products. The letters and codes used in Australia are unique because they were the first, they are compulsory and they reflect the relative risk of resistance evolving in each group. Since the introduction of mode of action labelling in Australia, other countries have adopted mode of action classification systems, however caution should be shown if cross-referencing mode of action between Australia and other countries, as many other countries use a different classification system.

The herbicide mode of action grouping and labelling system in Australia was revised in 2007. This is the first major revision of the classification system since its introduction. The original groupings were made several years ago based on limited knowledge about modes of action. Groupings have now been changed to improve the accuracy and completeness of the modes of action to ultimately enable more informed decisions to be made about herbicide rotation and resistance management. The general intent of groups based on their risk has not changed. However, six new herbicide mode of action groups were created to more accurately group herbicides. Titles describing some of the existing herbicide groups were also changed to more accurately describe those groups. The old groups E, F and K have the most changes. Most of the other groups remain unchanged. See details of the changes below.

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

### 3. CHANGES TO MODE OF ACTION GROUPINGS FOR SOME HERBICIDES

CropLife Australia revised the Australian herbicide mode of action groupings in 2007 and added six new groups (Groups H, O, P, Q, R and Z). The old groups E, F and K have the most changes (see table below). Most groups and herbicide products have not been changed, but the following active constituents and products have changed group and **the new group should be used when referring to the resistance management strategies**. Herbicide product registrants have three years (from February 2008) to update labels to reflect the new mode of action groups, but in the interim, some product labels may display the old mode of action group. **Where there is a temporary difference in mode of action group on labels, the new mode of action group in the table below should be used when choosing the appropriate resistance management strategy.**

<u>ACTIVE CONSTITUENT</u>	<u>PRODUCT NAME<sup>1</sup></u>	<u>CHEMICAL FAMILY</u>	<u>OLD MOA GROUP AND STRATEGY</u>	<u>NEW MOA GROUP AND STRATEGY</u>
Amitrole	Amitrole®, Illico®	Triazoles	GROUP F	GROUP Q
Asulam	Asulox®	Carbamates	GROUP K	GROUP R
bensulide	Prefar®	Phosphorodithioates	GROUP E	GROUP J
benzofenap	Taipan®, Viper®	Pyrazoles	GROUP F	GROUP H
clomazone	Command®, Viper®	Isoxazolidinones	GROUP F	GROUP Q
dichlobenil	Casoron®	Nitriles	GROUP K	GROUP O
Dithiopyr	Dimension®	Pyridines	GROUP E	GROUP D
DSMA	DSMA, Methar®	Organoarsenicals	GROUP K	GROUP Z
MSMA	Daconate®	Organoarsenicals	GROUP K	GROUP Z
Endothal	Endothal®	Dicarboxylic acids	GROUP K	GROUP Z
EPTC	Eptam®	Thiocarbamates	GROUP E	GROUP J
ethofumesate	Tramat®	Benzofurans	GROUP K	GROUP J
Flamprop	Mataven®	Arylamino propionic acids	GROUP K	GROUP Z
Isoxaben	Gallery®	Benzamides	GROUP K	GROUP O
isoxaflutole	Balance®	Isoxazoles	GROUP F	GROUP H
Molinate	Ordram®	Thiocarbamates	GROUP E	GROUP J
Naptalam	Alanap-L®	Phthalamates	GROUP K	GROUP P
Pebulate	Tillam®	Thiocarbamates	GROUP E	GROUP J
phenmedipham	Betanal®	Phenylcarbamates	GROUP K	GROUP C
propyzamide	Kerb®	Benzamides	GROUP K	GROUP D
Thiazopyr	Visor®	Pyridines	GROUP E	GROUP D
thiobencarb	Saturn®	Thiocarbamates	GROUP E	GROUP J
Triallate	Avadex®	Thiocarbamates	GROUP E	GROUP J
vernolate	Vernam®	Thiocarbamates	GROUP E	GROUP J

<sup>1</sup> The trade name of the first registered product or successor in each "Group" is listed under "Product Name". Refer to the website of the Australian Pesticides and Veterinary Medicines Authority (APVMA) ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products in the PUBCRIS database.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 4. HERBICIDES ARE GROUPED BY MODE OF ACTION AND RANKED BY RESISTANCE RISK

Growers and agronomists are now better aided to understand the huge array of herbicide products in the marketplace in terms of mode of action grouping and resistance risk by reference to the mode of action chart. All herbicide labels now carry the mode of action group clearly displayed such as: -

GROUP	G	HERBICIDE
-------	---	-----------

Know your herbicide groups to make use of this!

Not all mode of action groups carry the same risk for resistance development, therefore specific guidelines for Groups E, G, H, K, N, O, P and R have not been developed to date because there are no recorded cases of weeds resistant to members of these groups in Australia.

Products represented in Group A (mostly targeted at annual ryegrass and wild oats) and Group B (broadleaf and grass weeds) are HIGH RESISTANCE RISK herbicides and specific guidelines are written for use of these products in winter cropping systems.

Specific guidelines also follow for the MODERATE RESISTANCE RISK herbicides, Group C (annual ryegrass, wild radish and silver grass), Group D (annual ryegrass and fumitory), Group F (wild radish), Group I (wild radish and Indian hedge mustard), Group J (serrated tussock and giant Parramatta grass), Group L (barley grass, silver grass, square weed and capeweed), Group M (annual ryegrass, barnyard grass and liverseed grass) Group Q (annual ryegrass) and Group Z (wild oats).

Detailed programs for herbicide resistance management for weed control in rice are included (refer CropLife Australia website [www.croplifeaustralia.org.au](http://www.croplifeaustralia.org.au)). Details of herbicide resistance management plans in Liberty Link<sup>®</sup> Cotton, Roundup Ready<sup>®</sup> Cotton, Roundup Ready Flex<sup>®</sup> Cotton, Roundup Ready Canola<sup>®</sup> and Clearfield<sup>®</sup> Production Systems are available from Bayer, Monsanto and Nufarm respectively.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try to ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Make sure you rotate between products from different mode of action groups.

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 5. INTEGRATED WEED MANAGEMENT STRATEGIES

Strategies are designed to prevent and/or reduce the occurrence of resistance by adopting Integrated Weed Management (IWM) strategies. Do not rely on a single strategy to keep resistance at bay but integrate them into the crop production program. Some of the key strategies are: -

- Refer to specific guidelines for each herbicide mode of action group.
- Rotation of herbicide mode of action groups within and across years.
- **Keep accurate records of your herbicide applications on a paddock basis.**
- Read the herbicide product label and literature carefully and follow the instructions.
- Always use robust label rates
- Rotation of crop and variety.
- Identify and monitor your surviving weed populations and check for resistant weeds on your farm. Keep good records of weed populations.
- If a failure is suspected do not use the same product or product from the same mode of action group.
- Testing – confirm resistance exists.
- Control weed escape before the weeds set viable seed.
- Additional cultural weed control techniques to reduce seed banks, e.g. burning, cultivation, delayed sowing, competitive crops and varieties, green manuring, grazing and collection of weed seed at harvest.
- Do not introduce or spread weeds by contaminated seed, grain or hay.
- Consider crop and pasture topping.
- Attend training courses, e.g. GRDC IWM course, *ChemCert* and field days.
- Additional information can be obtained from: [CropLife Australia \(www.croplifeaustralia.org.au\)](http://www.croplifeaustralia.org.au), [Australian Glyphosate Sustainability Working Group \(www.glyphosateresistance.org.au\)](http://www.glyphosateresistance.org.au), [Grains Research & Development Corporation \(www.grdc.com.au\)](http://www.grdc.com.au) and State Government Departmental publications.
- Seek advice from local advisers (agronomists)

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is valid as at **15 September 2009**. All previous versions of this strategy are now invalid.  
Phone: 02 6230 6399 Email: [info@croplifeaustralia.org.au](mailto:info@croplifeaustralia.org.au) Fax: 02 6230 6355 Website: [www.croplifeaustralia.org.au](http://www.croplifeaustralia.org.au)

## 5. INTEGRATED WEED MANAGEMENT STRATEGIES (cont.)

### Weed control options for IWM

	<i>Herbicidal</i>	<i>Non-herbicidal</i>
<b>Crop phase</b>	Crop topping in pulse/legume crops Knockdown herbicides e.g. double knock strategy before sowing Selective herbicides before &/or after sowing – but ensure escapes don't set seed Utilising moderate resistance risk herbicides	Rotate crops Rotate varieties Grow a dense and competitive crop Cultivation Green/brown manure crops Delay sowing Cut crops for hay Burn stubbles / windrows Collect weed seeds at harvest
<b>Pasture phase</b>	Spray topping Winter cleaning Selective herbicides – but ensure escapes don't set seed	Good pasture competition Hay making or silage Cultivated fallow Grazing

Keep yourself informed and be pro-active in the fight-back against resistance. For further information on resistance management strategies, consult your reseller agronomist, farm consultant or Departmental Agronomist.

### **You can do something to reduce the impact!**

Follow the latest resistance management strategies described here.

#### **Note:**

In the specific guidelines for each mode of action group in the following pages, the boxes contain the chemical families, followed by a list of active constituents, with the trade name of the first registered product or successor in parentheses. The active constituents and some products that were moved to a different group in February 2008 are listed in bold in red.

For a complete list of registered products containing each active constituent, refer to the website of the Australian Pesticides and Veterinary Medicines Authority (APVMA) at [www.apvma.org.au](http://www.apvma.org.au) for the PUBCRIS database.

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 6. SPECIFIC GUIDELINES FOR GROUP A HERBICIDES

GROUP	A	HERBICIDE
-------	---	-----------

### High resistance risk

Group A resistance exists in Australia in the major grass weeds including annual ryegrass, wild oats, phalaris, brome grass, crab grass and barley grass. Resistance has developed in broadacre and vegetable situations.

Research has shown that on average as few as 6 applications to the same population of annual ryegrass can result in the selection of resistant individuals. A population can go from a small area of resistant individuals to a whole paddock failure in one season.

1. Fops, dims and dens are Group A herbicides and carry the same high resistance risk.
2. Where a Group A herbicide has been used on a particular paddock for control of any grass weed, avoid using a Group A herbicide to control the same grass weed in the following season, irrespective of the performance it gave.
3. Frequent application of Group A herbicides to dense weed populations is the worst case scenario for rapidly selecting for resistance.
4. Where resistance to a member of Group A is suspected or known to exist, there is a strong possibility of cross resistance to other Group A herbicides. Therefore use other control methods and herbicides of other mode of action groups in a future integrated approach.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try to ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Make sure you rotate between products from different mode of action groups.

GROUP A	Inhibitors of acetyl coA carboxylase (Inhibitors of fat synthesis/ACC'ase inhibitors)
<i>Aryloxyphenoxypropionates:</i> (Fops):	clodinafop (Topik®), cyhalofop (Barnstorm®), diclofop (Cheetah® Gold*, Decision®*, Hoegrass®, Tristar® Advance*), fenoxaprop (Cheetah® Gold*, Tristar® Advance*, Wildcat®), fluazifop (Fusilade®, Fusion®*), haloxyfop (Motsa®*, Verdict®), propaquizafop (Correct®), quizalofop (Targa®)
<i>Cyclohexanediones:</i> (Dims):	butoxydim (Falcon®, Fusion®*), clethodim (Motsa®*, Select®), profoxydim (Aura®), sethoxydim (Cheetah® Gold*, Decision®*, Sertin®), tepraloxydim (Aramo®), tralkoxydim (Achieve®)
<i>Phenylpyrazoles:</i> (Dens):	pinoxaden (Axial®)

\* **This product contains more than one active constituent**

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 7. SPECIFIC GUIDELINES FOR GROUP B HERBICIDES

GROUP	<b>B</b>	HERBICIDE
-------	----------	-----------

### High resistance risk

Group B resistance exists in Australia in the grass weeds, annual ryegrass, barley grass, brome grass, wild oats and crab grass and in at least 15 broadleaf weeds including: wild radish, common sowthistle, climbing buckwheat, turnip weed, wild mustard, Indian hedge mustard, prickly lettuce, wild turnip and African turnip weed. Resistance has developed in broadacre, rice and pasture situations.

Research has shown that on average as few as 4 applications to the same population of annual ryegrass can result in the selection of resistant individuals and as few as 6 applications for wild radish. A population can go from a small area of resistant individuals to a whole paddock failure in one season.

1. Avoid applying more than two Group B herbicides in any four year period on the same paddock.
2. **Broadleaf weed control:**  
If a pre-emergent application is made with a Group B herbicide for broadleaf weed control, monitor results and, if required, apply a follow up spray with a non-Group B herbicide for control of escapes and to reduce seed set.  
If a post-emergent application is made with a Group B herbicide for broadleaf weed control, this should preferably be as an APVMA approved tank-mix with another mode of action that controls or has significant activity against the target weed. If no APVMA approved tank-mix is available then monitor results and if required, apply a follow up spray with a non-Group B herbicide for control of escapes and to reduce seed set.  
A Group B herbicide may be used alone on flowering wild radish only if a Group B herbicide has not been previously used on that crop.
3. **Grass weed control:**  
If there are significant escapes following the herbicide application consider using another herbicide with a different mode of action or another control method to stop seed set.
4. **Clearfield Systems:**  
Where OnDuty®, Midas® and Intervix® are used refer to the Clearfield® Production System - Best Management Practice guide.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Make sure you rotate between products from different mode of action groups.

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

<b>GROUP B</b>	<b>Inhibitors of acetolactate synthase (ALS inhibitors)</b>
<i>Sulfonylureas:</i> <i>(SUs):</i>	azimsulfuron (Gulliver <sup>®</sup> ), bensulfuron (Londax <sup>®</sup> ), chlorsulfuron (Glean <sup>®</sup> ), ethoxysulfuron (Hero <sup>®</sup> ), halosulfuron (Sempra <sup>®</sup> ), iodosulfuron (Hussar <sup>®</sup> ), mesosulfuron (Atlantis <sup>®</sup> ), metsulfuron (Ally <sup>®</sup> , Harmony <sup>®</sup> M, Trounce <sup>®*</sup> ), rimsulfuron (Titus <sup>®</sup> ), sulfometuron (Oust <sup>®</sup> ), sulfosulfuron (Monza <sup>®</sup> ), thifensulfuron (Harmony <sup>®</sup> M), triasulfuron, (Logran <sup>®</sup> , Logran <sup>®</sup> B-Power <sup>®*</sup> ), tribenuron (Express <sup>®</sup> ), trifloxysulfuron (Envoke <sup>®</sup> , Krismat <sup>®*</sup> )
<i>Imidazolinones:</i> <i>(Imis):</i>	imazamox (Raptor <sup>®</sup> , Intervix <sup>®</sup> ), imazapic (Flame <sup>®</sup> , Midas <sup>®</sup> , OnDuty <sup>®</sup> ), imazapyr (Arsenal Xpress <sup>®</sup> , Midas <sup>®</sup> , OnDuty <sup>®</sup> , Intervix <sup>®</sup> , Lightning <sup>®</sup> ), imazethapyr (Spinnaker <sup>®</sup> , Lightning <sup>®</sup> )
<i>Triazolopyrimidines:</i> <i>(Sulfonamides):</i>	flumetsulam (Broadstrike <sup>®</sup> ), florasulam (Conclude <sup>®</sup> , Torpedo <sup>®</sup> , X-Pand <sup>®</sup> ), metosulam (Eclipse <sup>®</sup> ), pyroxsulam (Crusader <sup>®</sup> )
<i>Pyrimidinylthiobenzoates:</i>	pyrithiobac-Na (Staple <sup>®</sup> )

\* ***This product contains more than one active constituent***

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 8. SPECIFIC GUIDELINES FOR GROUP C HERBICIDES

GROUP	C	HERBICIDE
-------	---	-----------

### Moderate resistance risk

Group C resistance is known to exist in Australia in the weeds annual ryegrass, wild radish, liverseed grass, silver grass, stinging nettles and barnyard grass. Resistance has developed in broadacre, horticultural and non-crop situations.

CropLife Australia gives specific guidelines for the use of Group C herbicides in triazine tolerant canola and in winter legume crops, following increasing reports of resistance development: -

1. Avoid using Group C herbicides in the same paddock in consecutive years. Growing TT Canola in a paddock treated with triazine herbicides in the previous season is a high resistance risk and is not recommended.
2. **For use of triazines in triazine tolerant canola (TT): -**
  - Avoid dry sowing in heavily weed infested paddocks. Wait for a second weed germination after the opening rains in weedy paddock situations. Use a pre-plant knockdown or cultivation to maximise weed control at this stage.
  - Adapt the weed control program to the anticipated weed spectrum and pressure –
    - a) Broadleaf weeds and annual ryegrass  
Use simazine or atrazine or metribuzin plus trifluralin pre-emergence. A follow up with a Group A herbicide (if ryegrass is susceptible) or atrazine may be necessary
    - b) Broadleaf weeds only  
Use atrazine post-emergence
  - Watch and record for weed escapes, especially in paddocks with a long history of Group C use.

Consult the '**Integrated Weed Management Strategy for TT Canola**' for further details. (refer to manufacturing companies)

3. **The resistance status of the “at-risk” weeds should be determined prior to sowing.**  
The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Make sure you rotate between products from different mode of action groups.

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

CHEMICAL FAMILY	ACTIVE CONSTITUENT (FIRST REGISTERED TRADE NAME)
<b>GROUP C</b>	<b>Inhibitors of photosynthesis at photosystem II (PS II inhibitors)</b>
<i>Triazines:</i>	ametryn (Amigan <sup>®</sup> , Primatol Z <sup>®</sup> , Gesapax <sup>®</sup> Combi*, Krismat <sup>®</sup> ), atrazine (Gesaprim <sup>®</sup> , Gesapax <sup>®</sup> Combi*, Primextra <sup>®</sup> Gold*), cyanazine (Bladex <sup>®</sup> ), prometryn (Gesagard <sup>®</sup> , Cotogard <sup>®</sup> , Bandit <sup>®</sup> ), propazine (Agaprop <sup>®</sup> ), simazine (Gesatop <sup>®</sup> ), terbutryn (Amigan <sup>®</sup> , Igran <sup>®</sup> , Agtryne <sup>®</sup> MA*)
<i>Triazinones:</i>	hexazinone (Velpar <sup>®</sup> L, Velpar <sup>®</sup> K4*), metribuzin (Sencor <sup>®</sup> )
<i>Uracils:</i>	bromacil (Hyvar <sup>®</sup> , Krovar <sup>®*</sup> ), terbacil (Sinbar <sup>®</sup> )
<i>Pyridazinones:</i>	chloridazon (Pyramin <sup>®</sup> )
<i>Phenylcarbamates:</i>	<b>phenmedipham (Betanal<sup>®</sup>)</b>
<i>Ureas:</i>	diuron (Karmex <sup>®</sup> , Krovar <sup>®*</sup> , Velpar <sup>®</sup> K4*), fluometuron (Cotoran <sup>®</sup> , Cotogard <sup>®</sup> , Bandit <sup>®</sup> ), linuron (Afalon <sup>®</sup> ), methabenzthiazuron (Tribunil <sup>®</sup> ), siduron (Tupersan <sup>®</sup> ), tebuthiuron (Graslan <sup>®</sup> )
<i>Amides:</i>	propanil (Stam <sup>®</sup> )
<i>Nitriles:</i>	bromoxynil (Buctril <sup>®</sup> , Buctril <sup>®</sup> MA*, Barrel <sup>®*</sup> , Jaguar <sup>®*</sup> , Velocity <sup>®*</sup> ), ioxynil (Totril <sup>®</sup> , Actril DS*)
<i>Benzothiadiazinones:</i>	bentazone (Basagran <sup>®</sup> , Basagran <sup>®</sup> M60*)

**\* This product contains more than one active constituent**

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 9. SPECIFIC GUIDELINES FOR GROUP D HERBICIDES

<b>GROUP</b>	<b>D</b>	<b>HERBICIDE</b>
--------------	----------	------------------

### Moderate resistance risk

Resistance to Group D herbicides is known for an increasing number of populations of annual ryegrass and fumitory. Resistance has generally occurred after 10 -15 years of use of Group D herbicides.

Where possible, avoid the use of Group D herbicides on dense ryegrass populations. Consider using alternative methods of weed control to reduce weed numbers before applying herbicides.

Also consider using herbicides from other modes of action, where registrations permit.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Where possible, rotate between products from different mode of action groups.

<b>GROUP D</b>	<b>Inhibitors of microtubule assembly</b>
<i>Dinitroanilines:</i> <i>(DNAs):</i>	oryzalin (Surflan®, Yield®*), pendimethalin (Stomp®), trifluralin (Treflan®, Yield®*)
<i>Benzoic acids:</i>	chlorthal (Dacthal®, Prothal®*)
<i>Benzamides:</i>	<b>propyzamide (Kerb®)</b>
<i>Pyridines:</i>	<b>dithiopyr (Dimension®), thiazopyr (Visor®)</b>

\* ***This product contains more than one active constituent***

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 10. SPECIFIC GUIDELINES FOR GROUP F HERBICIDES

GROUP	F	HERBICIDE
-------	---	-----------

### Moderate resistance risk

Resistance to Group F herbicides is known for a small number of populations of wild radish. Resistance has generally occurred after a long history of use of Group F herbicides. The number of populations with Group F resistance is increasing following increased use of these herbicides.

Group F includes herbicides that reduce carotenoid biosynthesis through inhibition of phytoene desaturase (PDS).

Avoid applying Group F herbicides in any two consecutive years unless one application is a mixture with a different mode of action that is active on the same weed, or a follow up spray is conducted (using a different mode of action) to control escapes.

If applicable, apply a follow up spray with a non-Group F herbicide for control of escapes and to reduce seed set. In all cases, aim to ensure surviving weeds from any treatment do not set and shed viable seed.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Where possible, rotate between products from different mode of action groups.

GROUP F	<b>Bleachers: Inhibitors of carotenoid biosynthesis at the phytoene desaturase step (PDS inhibitors)</b>
<i>Nicotinilides:</i>	diflufenican (Brodal®, Jaguar®, Tigrex®, Chipco Spearhead®*)
<i>Picolinamides:</i>	picolinafen (Paragon®, Sniper®)
<i>Pyridazinones:</i>	norflurazon (Solicam®)

\* ***This product contains more than one active constituent***

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 11. SPECIFIC GUIDELINES FOR GROUP I HERBICIDES

GROUP	I	HERBICIDE
-------	---	-----------

### Moderate resistance risk

Resistance to Group I herbicides is known for a small number of populations of wild radish and Indian hedge mustard. Resistance has occurred after a long history of use of Group I herbicides. The number of populations with Group I resistance is increasing slowly.

Of particular concern is that apart from the resistance being in wild radish which is the most important broadleaf weed in broadacre agriculture, some populations are cross resistant to other modes of action e.g. Group F herbicides which can be important for control of wild radish in lupins where other selective non Group I options are limited. Because of the long soil life of wild radish seed, measures to reduce seed return to the soil would be useful for this weed. Wild radish seed that is confined to the top 5 cm soil has a shorter life than seed buried deeper.

As a general rule in high resistance risk situations

1. Avoid applying 2 applications of Group I herbicides alone onto the same population of weeds in the same season.
2. Where possible combine more than one mode of action in a single application. Each product should be applied at rates sufficient for control of the target weed alone to reduce the likelihood of weeds resistant to the Group I herbicide surviving.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Make sure you rotate between products from different mode of action groups.

GROUP I	Disruptors of plant cell growth
<i>Phenoxy</i> carboxylic acids: ( <i>Phenoxy</i> s):	2,4-D (Amicide <sup>®</sup> , Actril DS <sup>®*</sup> , <b>Pyresta<sup>®*</sup></b> ), 2,4-DB (Trifolamine <sup>®</sup> ), dichlorprop (Lantana 600 <sup>®</sup> ), MCPA (MCPA, Buctril <sup>®</sup> MA*, Banvel M <sup>®*</sup> , Conclude <sup>®</sup> , Midas <sup>®*</sup> , Paragon <sup>®*</sup> , Tigrex <sup>®*</sup> , Barrel <sup>®*</sup> , Tordon 242 <sup>®*</sup> , Basagran <sup>®</sup> M60*, Chipco Spearhead <sup>®*</sup> , Agtryne <sup>®</sup> MA*, Precept <sup>®*</sup> ), MCPB (Legumine <sup>®</sup> ), mecoprop (Mecopropamine <sup>®</sup> , Mecoban <sup>®</sup> , Methar Tri-Kombi <sup>®*</sup> )
<i>Benzoic</i> acids:	dicamba (Banvel <sup>®</sup> , Banvel M <sup>®*</sup> , Barrel <sup>®*</sup> , Mecoban <sup>®</sup> Methar Tri-Kombi <sup>®*</sup> )
<i>Pyridine</i> carboxylic acids: ( <i>Pyridines</i> ):	aminopyralid (Hotshot <sup>®*</sup> , Grazon Extra <sup>®*</sup> ), clopyralid (Lontrel <sup>®</sup> , Torpedo <sup>®*</sup> Chipco Spearhead <sup>®*</sup> ), fluroxypyr (Starane <sup>®</sup> , Hotshot <sup>®*</sup> ), picloram (Tordon <sup>®</sup> , Tordon 242 <sup>®*</sup> , Grazon <sup>®*</sup> , Grazon Extra <sup>®*</sup> ), triclopyr (Garlon <sup>®</sup> , Grazon <sup>®*</sup> , Grazon Extra <sup>®*</sup> )
<i>Quinoline</i> carboxylic acids:	quinclorac (Drive <sup>®</sup> )

**\* This product contains more than one active constituent**

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 12. SPECIFIC GUIDELINES FOR GROUP J HERBICIDES

GROUP	J	HERBICIDE
-------	---	-----------

### Moderate resistance risk

There are isolated cases of weeds resistant to Group J in Australia. There are 2 populations of serrated tussock and 6 populations of giant Parramatta grass resistant to flupropanate

To assist in delaying the onset of resistance, consider alternating with herbicides from other modes of action.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Where possible, rotate between products from different mode of action groups.

GROUP J	Inhibitors of fat synthesis (Not ACCase inhibitors)
Chlorocarbonic acids:	2,2-DPA (Dalapon®), flupropanate (Frenock®)
Thiocarbamates:	<b>EPTC (Eptam®), molinate (Ordram®), pebulate (Tillam®), prosulfocarb (Boxer® Gold*), thiobencarb (Saturn®), triallate (Avadex®), vernolate (Vernam®)</b>
Phosphorodithioates:	<b>bensulide (Prefar®)</b>
Benzofurans:	<b>ethofumesate (Tramat®)</b>

\* **This product contains more than one active constituent**

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 13. SPECIFIC GUIDELINES FOR GROUP L HERBICIDES

GROUP	L	HERBICIDE
-------	---	-----------

### Moderate resistance risk

Group L resistance exists in Australia in barley grass (2 species), silver grass, capeweed and square weed. Most instances have occurred in long-term lucerne stands treated regularly with a Group L herbicide but Group L resistant barley grass has also occurred in no-till situations.

The following factors are common to all cases of Group L resistance:

- A Group L herbicide is the major or only herbicide used
- A Group L herbicide has been used for 12 – 15 years or more: and
- There has been minimal or no soil disturbance following application.

The risk of resistance to Group L herbicides is higher in zero tillage broadacre cropping. Other high resistance risk situations include: orchards, vineyards or pure lucerne stands where frequent applications of a Group L herbicide are made each season, cultivation is not used and there is reliance on a Group L herbicide alone for weed control.

Below are strategies that address these high resistance risk situations to reduce the risk of Group L resistance developing.

### Zero Tillage

1. Rotate Group L herbicides with other knockdown herbicides with a different mode of action.
2. Consider utilising the double knock technique<sup>2</sup> where glyphosate is sprayed first followed within 1 - 5 days by a paraquat application. A full label rate for the weed size targeted should be used for the paraquat application for resistance management.
3. Consider occasional mechanical cultivation to aid weed control.

### Lucerne

1. If using a Group L herbicide for winter cleaning, where possible include another mode of action e.g. diuron (Group C).
2. Use alternative modes of action to selectively control grass and broadleaf weeds.
3. Rotate Group L herbicides with other knockdown herbicides with a different mode of action prior to sowing lucerne and prior to sowing future crops in that paddock.

<sup>2</sup> *The double knock technique is defined as using a full cut cultivation OR the full label rate of a paraquat-based product ( Group L) following the glyphosate ( Group M) knockdown application*

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

**Horticulture**

1. Rotate Group L herbicides with other knockdown herbicides with a different mode of action.
2. Where possible use residual herbicides (that are effective on the same weeds as the Group L herbicides) where applicable either alone or in mixture with Group L herbicides.
3. Where possible use alternative modes of action to selectively control grass and broadleaf weeds.
4. Consider using the double knock technique where glyphosate is sprayed followed within 1-5 days by a paraquat application. A full label rate for the weed size targeted should be used for the paraquat application for resistance management.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Always try to apply herbicides to the smallest weed density. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Make sure you rotate between products from different mode of action groups.

<b>GROUP L</b>	<b>Inhibitors of photosynthesis at photosystem I (PSI inhibitors)</b>
<i>Bipyridyls:</i>	diquat (Reglone®, Spray Seed®*), paraquat (Gramoxone®, Spray Seed®*, Alliance®*)

\* ***This product contains more than one active constituent***

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 14. SPECIFIC GUIDELINES FOR GROUP M HERBICIDES

GROUP	<b>M</b>	HERBICIDE
-------	----------	-----------

### Moderate resistance risk

Group M resistance occurs in Australia in annual ryegrass, awnless barnyard grass and liverseed grass.

Herbicide resistance to glyphosate was first discovered in annual ryegrass in Australia in 1996. Since that time, a number of new cases of glyphosate resistance in annual ryegrass, awnless barnyard grass and liverseed grass have been confirmed.

The following factors are common to all cases of Group M resistance:

- A Group M herbicide is the major or only herbicide used
- A Group M herbicide has been used for 12 – 15 years or more: and
- There has been minimal or no soil disturbance following application.

Given the very important role of glyphosate in Australian farming systems, the Australian agricultural industry has developed strategies for sustainable use of glyphosate. For more information refer to the Australian Glyphosate Sustainability Working Group website <http://www.glyphosateresistance.org.au>

All cases of glyphosate resistant weeds confirmed to date share three common factors:

- Intensive (year to year) use of glyphosate
- Lack of rotation of other herbicide modes of action
- Little or no tillage/cultivation following the application of glyphosate

A number of these cases of ryegrass resistance to glyphosate have occurred in horticultural and non-cropping situations (eg. firebreaks, fencelines, driveways, irrigation ditches), with the balance occurring in no-till broadacre cropping systems.

Given the demonstrated propensity of annual ryegrass to develop resistance to multiple herbicide classes, Integrated Weed Management principles should be incorporated wherever possible to minimise the risk of selecting for glyphosate resistant ryegrass. Strategies may include the use of cultivation, the double knock technique<sup>3\*</sup>, strategic herbicide rotation, grazing, baling etc.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Always try to apply herbicides to the smallest weed density. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Make sure you rotate between products from different mode of action groups.

<b>GROUP M</b>	<b>Inhibitors of EPSP synthase</b>
<i>Glycines:</i>	glyphosate (Roundup®, Trounce®, Illico®, Arsenal Xpress®)

\* ***This product contains more than one active constituent***

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

<sup>3</sup> ***The double knock technique is defined as using a full cut cultivation OR the full label rate of a paraquat-based product ( Group L) following the glyphosate ( Group M) knockdown application.***

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 15. SPECIFIC GUIDELINES FOR GROUP Q HERBICIDES

GROUP	Q	HERBICIDE
-------	---	-----------

### Moderate resistance risk

Group Q resistance exists in Australia in annual ryegrass resistant to amitrole. This has only occurred in 3 populations and this type of resistance is rare in Australia.

To assist in delaying the onset of resistance, consider alternating with herbicides from other modes of action eg. paraquat, glufosinate or glyphosate.

Consider using alternative methods of weed control to reduce weed numbers before applying herbicides.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Where possible, rotate between products from different mode of action groups.

GROUP Q	Bleachers: Inhibitors of carotenoid biosynthesis unknown target
Triazoles:	amitrole (Amitrole®, Illico®, Alliance®*)
Isoxazolidinones:	clomazone (Command®, Viper®*)

\* ***This product contains more than one active constituent***

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

## 16. SPECIFIC GUIDELINES FOR GROUP Z HERBICIDES

GROUP	Z	HERBICIDE
-------	---	-----------

### Moderate resistance risk

Group Z resistance exists in Australia in wild oats resistant to flamprop. Many of these flamprop resistant wild oats also show cross resistance to Group A herbicides.

To assist in delaying the onset of resistance, consider alternating with herbicides from other modes of action.

Consider using alternative methods of weed control to reduce weed numbers before applying herbicides. These may include summer crop rotations, delayed sowing to control wild oats with a knockdown herbicide, higher seeding rates, brown manuring to stop seed set, etc.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try and ensure surviving weeds from any treatment do not set and shed viable seed. Keep to the integrated strategies mentioned in this brochure including rotation of mode of action groups. Where possible, rotate between products from different mode of action groups.

GROUP Z	Herbicides with unknown and probably diverse sites of action
<i>Arylamino propionic acids:</i>	<b>flamprop (Mataven L®)</b>
<i>Dicarboxylic acids:</i>	<b>endothal (Endothal®)</b>
<i>Organoarsenicals:</i>	<b>DSMA (Methar®), MSMA (Daconate®)</b>

\* ***This product contains more than one active constituent***

List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)) to obtain a complete list of registered products from the PUBCRIS database. The active constituents and some products that were moved to a different mode of action group in February 2008 are listed in bold red type

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product data base for contemporary information on products and actives. The data base can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.